

Benlysta (Belimumab) IV Infusion Order Form

Full Name:	DOB:/
Phone:	Email:
Address:	
• Weight:	lb/kg
Allergies: □ NKE	DA 🗆
• Cu Treatment Sta	atus: New Continued Last Treatment://
Diagnosis (ICD-10 S	
M32.9 – Svstemic lupi	us erythematosus, unspecified
	I systemic lupus erythematosus
=	an or system involvement, unspecified
M32.11 – SLE with lun	•
	8 mvoivement
	_
M32.12 – SLE with kidi	ney involvement
M32.12 – SLE with kidi M32.13 – SLE with gas	ney involvement strointestinal involvement
M32.12 – SLE with kidi M32.13 – SLE with gas Other:	ney involvement
M32.12 – SLE with kidi M32.13 – SLE with gas	ney involvement strointestinal involvement
M32.12 – SLE with kidi M32.13 – SLE with gas Other: Infusion Order	ney involvement strointestinal involvement
M32.12 – SLE with kidi M32.13 – SLE with gas Other: Infusion Order • □ 10 mg/kg IV or	ney involvement strointestinal involvement ICD-10 Code:
M32.12 – SLE with kide M32.13 – SLE with gas Other: Infusion Order 10 mg/kg IV or 10 mg/kg IV or	ney involvement strointestinal involvement ICD-10 Code: nce every 2 weeks (initial 3 doses)
M32.12 – SLE with kidi M32.13 – SLE with gas Other: Infusion Order 10 mg/kg IV or 10 mg/kg IV or	ney involvement strointestinal involvement ICD-10 Code: nce every 2 weeks (initial 3 doses) nce every 4 weeks (maintenance)
M32.12 – SLE with kiding M32.13 – SLE with gas Other: Infusion Order 10 mg/kg IV or Infuse over approximate Refills:	ney involvement strointestinal involvement ICD-10 Code: nce every 2 weeks (initial 3 doses) nce every 4 weeks (maintenance)
M32.12 – SLE with kiding M32.13 – SLE with gas other: Infusion Order 10 mg/kg IV or 10 mg/kg I	ney involvement strointestinal involvement ICD-10 Code: nce every 2 weeks (initial 3 doses) nce every 4 weeks (maintenance)
M32.12 – SLE with kiding M32.13 – SLE with gas Other: Infusion Order 10 mg/kg IV or Infuse over approximate Refills: Line Use & Access	ney involvement strointestinal involvement ICD-10 Code: nce every 2 weeks (initial 3 doses) nce every 4 weeks (maintenance) roximately 1 hour
M32.12 – SLE with kiding M32.13 – SLE with gas of ther: Infusion Order Infusion Order Infusion Order Infusion Order Infusion Order Refills: Line Use & Access	ney involvement strointestinal involvement ICD-10 Code: nce every 2 weeks (initial 3 doses) nce every 4 weeks (maintenance) roximately 1 hour
M32.12 – SLE with kidi M32.13 – SLE with gas Other: Infusion Order 10 mg/kg IV of Infuse over appr Refills: Line Use & Access Start PIV — Access C	ney involvement strointestinal involvement ICD-10 Code: nce every 2 weeks (initial 3 doses) nce every 4 weeks (maintenance) roximately 1 hour CVC Use PICC Line Flush per standard infusion protocol

◆ <u>Premedication</u>
□ Acetaminophen: □ 500 mg □ 1000 mg □ PO
□ Diphenhydramine: □ 25 mg □ 50 mg □ PO □ IV
□ Solu-Medrol: □ 125 mg □ IV
•
Laboratory Monitoring
□ CBC □ CMP □ CRP □ ANA □ C3/C4 □ Other: □ Frequency: □ Each dose □ Monthly □ Other: □ Physician office will order labs only Clinical Documentation Checklist
□ Recent progress notes □ Last H&P □ Lab results □ Medication list
Ordering Provider & Demographics
• Name:
• NPI: License #:
• Contact: Phone: Fax:
• Email:
Signature: Date: / /