

Remicade (infliximab) IV Infusion Order Form

405-442-7577 📹 405-442-7223	
orders@premierinfusioncenter.com	
♣ Patient Information	
 Full Name: Date of Birth:// Phone: Email: 	
 Address: kg	
 Weight: kg	
Treatment Status: □ New □ Continued Last Treatment Date://	
 Diagnosis (ICD-10 Selection) 	
☐ K50.90 – Crohn's disease, unspecified	
☐ K51.90 – Ulcerative colitis, unspecified	
☐ M06.9 – Rheumatoid arthritis, unspecified	
☐ M45.9 – Ankylosing spondylitis	
☐ L40.52 – Psoriatic arthritis	
☐ L40.0 – Plaque psoriasis	
☐ Other: ICD-10 Code:	
• Infusion Order	
• Induction Phase:	
\square 5 mg/kg IV at Weeks 0, 2, and 6	
• Maintenance Phase:	
□ 5 mg/kg IV every 8 weeks	
☐ Alternative: mg/kg IV every weeks	
• \square Infuse over ≥ 2 hours (may reduce to 1 hour if tolerated)	
• ☐ Use 0.2-micron low-protein binding in-line filter	
• □ Observation: □ 30 minutes post-infusion	
• \square Refills: \square None \square 6 months \square 12 months \square Other:	

• Line Use &	Access			
☐ Start PIV protocol	□ Access CVC	☐ Use PICC Line	☐ Flush per s	standard infusion
• Adverse R	eaction & Anaphyl	axis Orders		
		l (premierinfusioncent tion orders to 405-442		
• Premedica	tion (Recommende	ed)		
☐ Diphenhydra ☐ Cetirizine: ☐ ☐ Methylpredn	\mathcal{C}	□ PO □ 50 mg □ PO □ IV Dose:		
Laboratory				_
☐ Hepatitis B I☐ Other:	g (within 12 months Panel (within 3 years		□ Monthly	□ Other:
☐ Physician of	fice will order labs of	only		
Clinical Do□ Recent progr	ocumentation Chec	st H&P □ Lab res	sults □ Medi	ecation list
	-			
	Provider & Demog	-		
Name:NPI:ContactEmail:	Licen	se #: Phone:		Fax:
				/